

Mouse Anti Human CD11b Purified

PRODUCT INFORMATION

CLONE: HI11b
ISOTYPE: Mouse IgG2b
CATALOG#: A6241/A6241-bulk
CONTENTS: Purified antibody buffered in 10mM PBS (PH 7.0) with 0.1% NaN₃.

DESCRIPTION

CD11b McAb recognizes the 165 KD integrin α M-chain associated with CD18 (95KD integrin β 2) forming heterodimer glycoprotein called CD11b/CD18 (Mac-1). Leucocytes express four CD11/CD18 integrins which comprise a common β 2 subunit and distinctive α subunits. The other three integrins are known as CD11a/CD18, CD11c/CD18 and CD11d/CD18. Expression of the CD11b subunit on the leucocyte cell surface requires the presence of the CD18 subunit. Patients who have mutated CD18 genes do not express cell membrane Mac-1 (or the other β 2 integrins), a condition known as leucocyte adhesion deficiency 1 (LAD-1), in which patients have major problems with bacterial infections. Mac-1 is most highly expressed on neutrophils and monocytes/macrophages, a subset of NK cells, and activated lymphocytes. Mac-1 operates as a phagocytic receptor, and has a role in myeloid cell transmigration and is also as a receptor for iC3b, CD54 (ICAM-1), CD102 (ICAM-2) and fibrinogen.

PREPARATION

The purified reagent is effective for indirect immunofluorescence staining of human cells for flow cytometric analysis and is tested for immunohistochemical staining of acetone-fixed frozen sections and formalin-fixed paraffin sections.

USAGE

The purified reagent is effective for indirect immunofluorescence staining of human cells for flow cytometric analysis and is tested for immunohistochemical staining of acetone-fixed frozen sections.

STORAGE

Store at 4°C. For long-term storage aliquot and store at -20°C. Avoid freeze and thaw cycles.

Use For Research Only