



## Mouse Anti Human CD2 FITC

### PRODUCT INFORMATION

**CLONE:** HIT11  
**ISOTYPE:** Mouse IgG1  
**WS.No.:** V 5T CD02.04  
**CATALOG#:** A6042/A6052  
**CONTENTS:** FITC - conjugated antibody buffered in 10mM PBS (PH 7.0) with 0.05% NaN<sub>3</sub> and 1% BSA.

### DESCRIPTION

CD2 McAb recognizes a 50 KD type I single-chain transmembrane glycoprotein called LFA-2 (lymphocyte function associated antigen-2) or sheep red blood cell (SRBC) receptor. CD2 antigen is expressed on about 70%-80% of peripheral blood lymphocytes, 80%-90% of thymocytes and NK cells. The primary ligand for CD2 is CD58 (LFA-3). The studies in previous and recent have revealed that CD48, CD59 and CD15 also act as ligands for CD2. CD2 antigen plays a role in T cell signaling and in lymphocyte adhesion and is an alternative pathway of T cell activation. There are membrane form on cell surface and soluble form in serum in CD2 antigen.

### PREPARATION

The monoclonal antibody is purified from ascites by protein G affinity chromatography and is conjugated with FITC under optimum conditions.

### USAGE

The FITC conjugation is tested for flow cytometric analysis using 20 $\mu$ l/10<sup>6</sup> cells or 100 $\mu$ l peripheral blood cells.

### STORAGE

Store at 4°C, should not be frozen and avoid prolonged exposure to light.

### REFERENCES

Schlossman, S., L. Bloumsell, W. Gilks, et al., eds. 1995. Leucocyte Typing V: White Cell Differentiation Antigens. P: 245, 346-348 Oxford University Press, New York.

Use For Research Only