



Mouse Anti Human CD16 PE

PRODUCT INFORMATION

CLONE: HI16a
ISOTYPE: Mouse IgG1, κ
CATALOG#: A6323/A6333
CONTENTS: R-PE conjugated antibody buffered in 10mM PBS (pH 7.0) with 0.05% NaN₃ and 1% BSA.

DESCRIPTION

CD16 McAb recognizes a 50-60KD transmembrane molecule, low affinity FC receptor for IgG (Fc γ RIII), which is a NK cell-associated antigen. CD16 antigen is expressed on the large majority of NK cells, granulocytes (but on eosinophils), monocytes/macrophages and a subset of T cells. CD16 antigen may involve in signal transduction, NK cell activation and antibody-dependent cellular cytotoxicity (ADCC).

PREPARATION

The monoclonal antibody is purified from ascites by protein G affinity chromatography and is conjugated with R-PE under optimum conditions.

USAGE

The R-PE conjugation is tested for flow cytometric analysis using 20 μ l/10⁶ cells or 100 μ l peripheral blood cells.

STORAGE

Store at 4°C, should not be frozen and avoid prolonged exposure to light.

REFERENCES

1. Knapp, W., B. Dorken, E. P. Rieber, et al., eds. 1989. Leucocyte Typing IV: White Cell Differentiation Antigens. P:582 Oxford University Press, New York.
2. Barclay, N.A., M.L.Birkeland, M.H. Brown, et al., eds. 1993. The Leucocyte Antigen FactsBook, CD16 Section, Academic Press Inc., San Diego, California, p.136.

Use For Research Only